



# Search for $\Lambda_c^+ \rightarrow pK^+\pi^-$ and $D_s^+ \rightarrow K^+K^+\pi^-$ using genetic programming event selection

FOCUS Collaboration <sup>1</sup>

J.M. Link <sup>a</sup>, P.M. Yager <sup>a</sup>, J.C. Anjos <sup>b</sup>, I. Bediaga <sup>b</sup>, C. Castromonte <sup>b</sup>, A.A. Machado <sup>b</sup>,  
J. Magnin <sup>b</sup>, A. Massafferri <sup>b</sup>, J.M. de Miranda <sup>b</sup>, I.M. Pepe <sup>b</sup>, E. Polycarpo <sup>b</sup>,  
A.C. dos Reis <sup>b</sup>, S. Carrillo <sup>c</sup>, E. Casimiro <sup>c</sup>, E. Cuautle <sup>c</sup>, A. Sánchez-Hernández <sup>c</sup>,  
C. Uribe <sup>c</sup>, F. Vázquez <sup>c</sup>, L. Agostino <sup>d</sup>, L. Cinquini <sup>d</sup>, J.P. Cumalat <sup>d</sup>, B. O'Reilly <sup>d</sup>,  
I. Segoni <sup>d</sup>, K. Stenson <sup>d</sup>, J.N. Butler <sup>e</sup>, H.W.K. Cheung <sup>e</sup>, G. Chiodini <sup>e</sup>, I. Gaines <sup>e</sup>,  
P.H. Garbincius <sup>e</sup>, L.A. Garren <sup>e</sup>, E. Gottschalk <sup>e</sup>, P.H. Kasper <sup>e</sup>, A.E. Kreymer <sup>e</sup>,  
R. Kutschke <sup>e</sup>, M. Wang <sup>e</sup>, L. Benussi <sup>f</sup>, M. Bertani <sup>f</sup>, S. Bianco <sup>f</sup>, F.L. Fabbri <sup>f</sup>,  
S. Pacetti <sup>f</sup>, A. Zallo <sup>f</sup>, M. Reyes <sup>g</sup>, C. Cawfield <sup>h</sup>, D.Y. Kim <sup>h</sup>, A. Rahimi <sup>h</sup>, J. Wiss <sup>h</sup>,  
R. Gardner <sup>i</sup>, A. Kryemadhi <sup>i</sup>, Y.S. Chung <sup>j</sup>, J.S. Kang <sup>j</sup>, B.R. Ko <sup>j</sup>, J.W. Kwak <sup>j</sup>,  
K.B. Lee <sup>j</sup>, K. Cho <sup>k</sup>, H. Park <sup>k</sup>, G. Alimonti <sup>1</sup>, S. Barberis <sup>1</sup>, M. Boschini <sup>1</sup>, A. Cerutti <sup>1</sup>,  
P. D'Angelo <sup>1</sup>, M. DiCorato <sup>1</sup>, P. Dini <sup>1</sup>, L. Edera <sup>1</sup>, S. Erba <sup>1</sup>, P. Inzani <sup>1</sup>, F. Leveraro <sup>1</sup>,  
S. Malvezzi <sup>1</sup>, D. Menasce <sup>1</sup>, M. Mezzadri <sup>1</sup>, L. Moroni <sup>1</sup>, D. Pedrini <sup>1</sup>, C. Pontoglio <sup>1</sup>,  
F. Prelz <sup>1</sup>, M. Rovere <sup>1</sup>, S. Sala <sup>1</sup>, T.F. Davenport III <sup>m</sup>, V. Arena <sup>n</sup>, G. Boca <sup>n</sup>, G. Bonomi <sup>n</sup>,  
G. Gianini <sup>n</sup>, G. Liguori <sup>n</sup>, D. Lopes Pegna <sup>n</sup>, M.M. Merlo <sup>n</sup>, D. Pantea <sup>n</sup>, S.P. Ratti <sup>n</sup>,  
C. Riccardi <sup>n</sup>, P. Vitulo <sup>n</sup>, C. Göbel <sup>o</sup>, H. Hernandez <sup>p</sup>, A.M. Lopez <sup>p</sup>, H. Mendez <sup>p</sup>,  
A. Paris <sup>p</sup>, J. Quinones <sup>p</sup>, J.E. Ramirez <sup>p</sup>, Y. Zhang <sup>p</sup>, J.R. Wilson <sup>q</sup>, T. Handler <sup>r</sup>,  
R. Mitchell <sup>r</sup>, D. Engh <sup>s</sup>, M. Hosack <sup>s</sup>, W.E. Johns <sup>s</sup>, E. Luiggi <sup>s</sup>, J.E. Moore <sup>s</sup>,  
M. Nehring <sup>s</sup>, P.D. Sheldon <sup>s</sup>, E.W. Vaandering <sup>s</sup>, M. Webster <sup>s</sup>, M. Sheaff <sup>t</sup>

<sup>a</sup> University of California, Davis, CA 95616, USA

<sup>b</sup> Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, RJ, Brazil

<sup>c</sup> CINVESTAV, 07000 México City, DF, Mexico

<sup>d</sup> University of Colorado, Boulder, CO 80309, USA

<sup>e</sup> Fermi National Accelerator Laboratory, Batavia, IL 60510, USA

<sup>f</sup> Laboratori Nazionali di Frascati dell'INFN, Frascati I-00044, Italy

<sup>g</sup> University of Guanajuato, 37150 Leon, Guanajuato, Mexico

<sup>h</sup> University of Illinois, Urbana-Champaign, IL 61801, USA

<sup>i</sup> Indiana University, Bloomington, IN 47405, USA

<sup>j</sup> Korea University, Seoul 136-701, South Korea

<sup>k</sup> Kyungpook National University, Taegu 702-701, South Korea

<sup>1</sup> INFN and University of Milano, Milano, Italy<sup>m</sup> University of North Carolina, Asheville, NC 28804, USA<sup>n</sup> Dipartimento di Fisica Nucleare e Teorica and INFN, Pavia, Italy<sup>o</sup> Pontifícia Universidade Católica, Rio de Janeiro, RJ, Brazil<sup>p</sup> University of Puerto Rico, Mayaguez, PR 00681, USA<sup>q</sup> University of South Carolina, Columbia, SC 29208, USA<sup>r</sup> University of Tennessee, Knoxville, TN 37996, USA<sup>s</sup> Vanderbilt University, Nashville, TN 37235, USA<sup>t</sup> University of Wisconsin, Madison, WI 53706, USA

Received 28 July 2005; received in revised form 10 August 2005; accepted 11 August 2005

Available online 22 August 2005

Editor: L. Rolandi

---

**Abstract**

We apply a genetic programming technique to search for the doubly Cabibbo suppressed decays  $\Lambda_c^+ \rightarrow pK^+\pi^-$  and  $D_s^+ \rightarrow K^+K^+\pi^-$ . We normalize these decays to their Cabibbo favored partners and find  $\text{BR}(\Lambda_c^+ \rightarrow pK^+\pi^-)/\text{BR}(\Lambda_c^+ \rightarrow pK^-\pi^+) = (0.05 \pm 0.26 \pm 0.02)\%$  and  $\text{BR}(D_s^+ \rightarrow K^+K^+\pi^-)/\text{BR}(D_s^+ \rightarrow K^-K^+\pi^+) = (0.52 \pm 0.17 \pm 0.11)\%$  where the first errors are statistical and the second are systematic. Expressed as 90% confidence levels (CL), we find  $< 0.46$  and  $< 0.78\%$ , respectively. This is the first successful use of genetic programming in a high energy physics data analysis.

© 2005 Elsevier B.V. Open access under [CC BY license](#).

PACS: 13.25.Ft; 13.30.Eg

Keywords: Genetic programming

---

Cabibbo suppressed (CS) and doubly Cabibbo suppressed (DCS) decays are important in helping us understand the dynamics of hadronic decay processes. DCS decays are unique to the charmed hadrons; charm is the only heavy up-type quark that hadronizes. DCS decay rates are such that only DCS decays of  $D^+$  and  $D^0$  have been observed, while CS decays of nearly all the charmed hadrons have been observed. This Letter presents a search for DCS decays of  $\Lambda_c^+$  and  $D_s^+$ . Both branching ratios are expected to be small. Naïve expectations place DCS branching ratios around  $\tan^4 \theta_c$ , or about 0.25%, relative to their Cabibbo favored (CF) counterparts. Lipkin argues [1] that exact SU(3) symmetry would require the product of the DCS relative branching ratios  $\text{BR}(D^+ \rightarrow K^+\pi^-\pi^+)/\text{BR}(D^+ \rightarrow K^-\pi^+\pi^+)$  and  $\text{BR}(D_s^+ \rightarrow K^+K^+\pi^-)/\text{BR}(D_s^+ \rightarrow K^-K^+\pi^-)$  to be exactly  $\tan^8 \theta_c$ . This means the latter should be about 0.07%; a much larger value requires

a large violation of flavor SU(3). In the  $\Lambda_c^+$  case, the CF normalizing mode has a  $c$ - $d$   $W^+$  exchange decay channel available, while the DCS decay mode may only proceed through spectator decays. The lifetime difference between  $\Lambda_c^+$  and  $\Xi_c^+$  shows us that this exchange mode is important, so we expect that the branching ratio for  $\Lambda_c^+ \rightarrow pK^+\pi^-$  should also be less than  $\tan^4 \theta_c$ .

We have applied a genetic programming (GP) [2] technique to search for the DCS decays  $D_s^+ \rightarrow K^+K^+\pi^-$  and  $\Lambda_c^+ \rightarrow pK^+\pi^-$  (charge-conjugate states are implied), neither of which have been observed. GP is a machine learning technique which evolves populations of programs (event filters in our case) over a series of generations. The genetic programming learning mechanism is modeled on biological and evolutionary principals and differs from some other machine learning solutions in that the form of the solution is not specified in advance but is determined by the complexity of the problem. We have found that this unbiased method can be more effective in separating signal events from backgrounds than standard,

---

E-mail address: [ewv@fnal.gov](mailto:ewv@fnal.gov) (E.W. Vaandering).

<sup>1</sup> See <http://www-focus.fnal.gov/authors.html> for additional author information.

cut based, methods. A full demonstration of this technique on the observed DCS decay  $D^+ \rightarrow K^+\pi^+\pi^-$  is given in Ref. [3].

These results use data taken with the charm photoproduction experiment FOCUS (FNAL-E831), an upgraded version of FNAL-E687 [4] which collected data using the Wideband photon beamline during the 1996–1997 Fermilab fixed-target run. The FOCUS experiment utilizes a forward multiparticle spectrometer to study charmed particles produced by the interaction of high energy photons ( $\langle E \rangle \approx 180$  GeV) [5] with a segmented BeO target. Charged particles are tracked within the spectrometer by two silicon microvertex detector systems. One system is interleaved with the target segments [6]; the other is downstream of the target region. These detectors provide excellent separation of the production and decay vertices. Further downstream, charged particles are tracked and momentum analyzed with a system of five multiwire proportional chambers [7] and two dipole magnets of opposite polarity. Three multicell threshold Čerenkov detectors are used to identify electrons, pions, kaons, and protons. FOCUS also contains a complement of hadronic and electromagnetic calorimeters and muon detectors.

We use loose analysis cuts on both DCS and CF decay modes to select initial samples of events for optimization by GP. The FOCUS Čerenkov algorithm [8] returns negative  $2 \times \log$ -likelihood values  $W_i(j)$  for

particle  $j$  and hypothesis  $i \in e, \pi, K, p$ . Differences between log-likelihoods are used as particle ID, such as  $\Delta W_{Kp}(p) \equiv W_K(p) - W_p(p)$  for “proton favored over kaon.” We require  $\Delta W_{\pi K}(K) > 2$  for all kaons in both decay modes. For protons from  $\Lambda_c^+$  candidates, we require  $\Delta W_{\pi p}(p) > 4$  and  $\Delta W_{Kp}(p) > 0$  in the initial selection. For the  $\Lambda_c^+$ , we also require that the separation between the production and decay vertices,  $\ell$ , is greater than 3 times its error,  $\sigma_\ell$ . For the  $D_s^+$ , the vertex separation requirement is  $\ell/\sigma_\ell > 6$ . For both  $\Lambda_c^+$  and  $D_s^+$ , the three decay fragments must form a vertex with a confidence level (CL)  $> 1\%$ , and a production vertex is formed by adding as many remaining tracks to the charm candidate as possible while maintaining a vertex CL  $> 1\%$ . One additional requirement is placed on the CF (DCS)  $D_s^+$  candidates: the  $K^-K^+\pi^+$  ( $K^+K^+\pi^-$ ) combination is rejected if, reconstructed as  $K^-\pi^+\pi^+$  ( $K^+\pi^+\pi^-$ ), the mass is within  $2\sigma$  of the nominal  $D^+$  mass. This cut removes a prominent reflection from the CF candidates and stabilizes the many fits done during the optimization process; it is applied to the DCS mode for consistency. The initial samples of  $\Lambda_c^+$  and  $D_s^+$  candidates in CF and DCS decay modes are shown in Fig. 1.

A GP framework (GPF) evolves and tests event filters. For each filter, we define a fitness

$$f \propto \frac{B_{\text{DCS}}}{S_{\text{CF}}^2} \times (1 + 0.005 \times N_{\text{nodes}}), \quad (1)$$




Fig. 1. Initial  $\Lambda_c^+$  (left) and  $D_s^+$  (right) data samples. The upper distributions show the CF decay candidates, the lower distributions show the DCS candidates. In the  $D_s^+$  plot, the  $D^+ \rightarrow K^-\pi^+\pi^+$  rejection cut described in the text is *not* applied and a contribution to the fit for this reflection is visible to the right of the  $D_s^+$  peak.




Fig. 2.  $\Lambda_c^+$  samples after selection. On the left is the CF normalizing mode, on the right, the remaining  $\Lambda_c^+ \rightarrow pK^+\pi^-$  candidates.

where  $B_{DCS}$  is the number of background events found in a fit of the DCS mass distribution which excludes the signal region, and  $S_{CF}$  is the fitted CF yield.  $S_{CF}/\sqrt{B_{DCS}}$  is proportional to the projected DCS significance assuming no real DCS events and equal CF and DCS selection efficiencies; squaring this quantity further emphasizes “better” filters and inverting it allows small fitnesses to describe good event filters.  $N_{nodes}$  is the total number of variables, constants, functions, and operators used in the filter and is included as a penalty term to encourage smaller filters and to attempt to eliminate the addition of nodes which do not select events based on physics.

For the data samples in Fig. 1, half the events (as explained later) along with a large number of variables (37 for  $\Lambda_c^+$ , 34 for  $D_s^+$ ), operators and functions (21), and constants are used as inputs to the GPF which randomly generates a large number of filters and calculates the fitness of each. The GPF preferentially selects filters for which this fitness is small to participate in breeding subsequent generations of filters. In this way subsequent generations develop filters with better average fitnesses. At the end of the process we use the filter with the single best fitness to select events for further analysis.

The variables and resulting filter used in the CF and DCS decays are identical. All variables commonly used in FOCUS analyses and some additional variables are allowed to be used in the event filter. These can be roughly broken into categories of vertexing, track quality, particle identification, production

and decay kinematics, away-side charm tagging and, for the  $\Lambda_c^+$ , evidence for decays of the excited states  $\Sigma_c^{(*)0,++}$ . A description of the variables<sup>2</sup> used, examples of the event filters constructed, and how the population of filters evolves over many generations can be found in Ref. [3]. In both cases we use 20 sub-populations of 1500 event filters per generation as described in Ref. [3].

When searching for  $\Lambda_c^+ \rightarrow pK^+\pi^-$  and  $D_s^+ \rightarrow K^+K^+\pi^-$  decays (with the signal regions masked), the GPF is allowed to run for 80 generations. The process is terminated when no improvement in fitness is observed for about 10 generations. The best  $\Lambda_c^+ \rightarrow pK^+\pi^-$  filter found has 45 nodes and uses 12 unique physics variables. The events selected are shown in Fig. 2. One can see that about 15% of the signal is retained compared to Fig. 1 while the backgrounds are reduced by a factor of  $\sim 1000$ . The distributions in both the CF and DCS cases are fit with a second degree polynomial<sup>3</sup> and a single Gaussian. In the DCS case, the Gaussian mean and  $\sigma$  are fixed to the CF values and we find  $1.2 \pm 6.6$  events. Correcting

<sup>2</sup> In addition to the variables described in Ref. [3], we add three additional variables:  $\Delta W_{K\pi}(\pi)$ , the number of tracks in the production vertex, and a value indicating if any of the vertex detector track segments are shared between two tracks.

<sup>3</sup> No significant reflections in  $\Lambda_c^+ \rightarrow pK^+\pi^-$  or  $D_s^+ \rightarrow K^+K^+\pi^-$  are seen in high-statistics MC studies (which include all known  $c\bar{c}$  decay processes) of these decays, so we are justified in using simple background shapes.




Fig. 3.  $D_s^+$  samples after selection. On the left is the CF normalizing mode, on the right, the remaining  $D_s^+ \rightarrow K^+ K^+ \pi^-$  candidates.

for the relative efficiency  $\epsilon_{CF}/\epsilon_{DCS} = 1.204 \pm 0.007$  (stat.) calculated with Monte Carlo (MC) simulations, we obtain a relative BR of

$$\frac{\text{BR}(\Lambda_c^+ \rightarrow p K^+ \pi^-)}{\text{BR}(\Lambda_c^+ \rightarrow p K^- \pi^+)} = (0.05 \pm 0.26)\%, \quad (2)$$

which is consistent with zero.

The best  $D_s^+ \rightarrow K^+ K^+ \pi^-$  filter found has 85 nodes and uses 15 unique physics variables. The events selected are shown in Fig. 3. The fits shown are performed identically to the  $\Lambda_c^+$  case except that an additional Gaussian is added to the CF distribution for the CS decay  $D^+ \rightarrow K^- K^+ \pi^+$ . We find  $27.5 \pm 9.2$  events in the DCS distribution which, corrected by the relative efficiency  $\epsilon_{CF}/\epsilon_{DCS} = 1.154 \pm 0.005$ , gives a relative BR of

$$\frac{\text{BR}(D_s^+ \rightarrow K^+ K^+ \pi^-)}{\text{BR}(D_s^+ \rightarrow K^- K^+ \pi^+)} = (0.52 \pm 0.17)\%. \quad (3)$$

In both cases our central values are calculated assuming non-resonant decays for the DCS case and the best known resonance models for the CF decays (the PDG [9] model for  $\Lambda_c^+$  and a FOCUS model for  $D_s^+$ ) as explained below.

To convert these relative BRs into upper limits including systematic errors, we use a method proposed by Convery [10] for incorporating systemic uncertainties on reconstruction efficiencies into BR measurements when a fit, rather than event counting, is used. In this case, the probability  $P(B)$  of the true BR being

$B$  is given by

$$p(B) \propto \frac{1}{\sqrt{\frac{B^2}{2\sigma_B^2} + \frac{\hat{S}^2}{2\sigma_S^2}}} \exp\left[\frac{-(B - \hat{B})^2}{2\left(\frac{B^2\sigma_S^2}{\hat{S}^2} + \sigma_B^2\right)}\right], \quad (4)$$

where  $\hat{B}$  is the fitted BR,  $\sigma_B$  is its error and  $\sigma_S/\hat{S}$  is the percent systematic error on the efficiency.  $P(B)$  is numerically integrated until the point at which 90% of the physical ( $\text{BR} > 0$ ) area is included. This point is reported as the 90% confidence level. If  $\sigma_S/\hat{S} \gg 10\%$ , this distribution has a long high-end tail, raising the 90% limit considerably.

We consider four sources of systematic error on our knowledge of the relative efficiencies of the CF and DCS decay modes. First, and negligible, is the number of MC events used. Second and third, we consider the effects of different resonance models for the DCS and CF states, respectively. Finally, we consider whether the evolved event selector may have different efficiencies for the CF and DCS modes.

In studying possible resonances for  $\Lambda_c^+ \rightarrow p K^+ \pi^-$  candidates, we calculate efficiencies as if the final state is entirely non-resonant or entirely  $\Lambda_c^+ \rightarrow \Delta(1232)^0 K^+$  or  $\Lambda_c^+ \rightarrow p K^*(892)^0$ . The systematic error is taken as the standard deviation of the three possible efficiencies. For the  $D_s^+ \rightarrow K^+ K^+ \pi^-$  candidates, we consider non-resonant decays,  $D_s^+ \rightarrow K^*(892)^0 K^+$ , and  $D_s^+ \rightarrow K_0^*(1430)^0 K^+$  in the same way. From these studies we find 5.3 and 10.7% sys-




Fig. 4. Relative BR limit determination for  $\Lambda_c^+ \rightarrow pK^+\pi^-$  (left) and  $D_s^+ \rightarrow K^+K^+\pi^-$  (right). The curves show the branching ratio probability for DCS decays relative to CF decays. The vertical axes are arbitrary. The shaded areas show the 90% integrals over the physical range.

tematic uncertainties on the  $\Lambda_c^+$  and  $D_s^+$  efficiencies, respectively.

The resonant structures of the CF decays are reasonably well known. For the  $\Lambda_c^+$ , we use two models, one from the PDG and another which excludes the  $\Lambda(1520)^0\pi^+$  decay mode. For the  $D_s^+$ , we consider an incoherent model based on the PDG averages and a coherent model [11] developed from the FOCUS data. From these studies we find 2.1 and 2.6% uncertainties on the  $\Lambda_c^+ \rightarrow pK^-\pi^+$  and  $D_s^+ \rightarrow K^-K^+\pi^+$  efficiencies, respectively.

Our final systematic contribution is motivated by the concern that the efficiency of the final GP-generated event filter may differ for the CF and DCS modes (after correction for kinematic acceptance of different final states) in a way that is not well modeled by MC. Since this is impossible to measure, we adopt a more rigorous test. We test if the event filter has the same efficiency on CF data and MC events. We do this by comparing the CF yields of data and MC samples before and after the event filter is applied.<sup>4</sup> For the  $\Lambda_c^+$ , we find that the event filter retains  $14.5 \pm 0.4\%$  and  $14.9 \pm 0.1\%$  of the data and MC events, respectively. For the  $D_s^+$ , we determine these quantities to be  $21.0 \pm 0.4\%$  and  $20.3 \pm 0.1\%$ . We take the differences between these numbers (neglecting the errors) as sys-

<sup>4</sup> The  $D^+ \rightarrow K^-\pi^+\pi^+$  rejection cut is applied with the event filter in the  $D_s^+$  case.

Table 1

Summary of systematic uncertainties. Listed are the percent uncertainties on the relative efficiencies of the DCS and CF decay modes from various sources

Source	Syst. unc. (%)	
	$\Lambda_c^+$	$D_s^+$
MC statistics	0.6	0.4
DCS resonances	5.3	10.7
CF resonances	2.1	2.6
GP filter	2.6	3.5
Total	6.3	11.6

tematic uncertainties; these cause 2.6 and 3.5% uncertainties on the relative efficiencies for  $\Lambda_c^+ \rightarrow pK^+\pi^-$  and  $D_s^+ \rightarrow K^+K^+\pi^-$ , respectively. All systematic uncertainties on the relative efficiencies are summarized in Table 1.

Finally, as mentioned above, we only use half (even-numbered) of the events in the optimization of the event filter. The final values use the event filter applied to all the events, but as a check, we divide the sample into events the GPF used and did not use. We measure the BR independently for these two samples and see no significant evidence for a difference, strongly suggesting that the GPF is not arbitrarily selecting or rejecting small numbers of events to artificially reduce backgrounds or enhance signals.

Using the total percent errors in Table 1 as  $\sigma_S/\hat{S}$  and the above BRs, statistical errors, and percent sys-

tematic errors, we integrate  $P(B)$  from Eq. (4) as described and find

$$\frac{\text{BR}(\Lambda_c^+ \rightarrow pK^+\pi^-)}{\text{BR}(\Lambda_c^+ \rightarrow pK^-\pi^+)} < 0.46\% \quad (5)$$

and

$$\frac{\text{BR}(D_s^+ \rightarrow K^+K^+\pi^-)}{\text{BR}(D_s^+ \rightarrow K^-K^+\pi^+)} < 0.78\%, \quad (6)$$

where the limits are at the 90% CL. We also determine effective systematic uncertainties for our measurements by calculating the uncertainty necessary, when added in quadrature to the statistical uncertainty, to cover the central 68% of the distribution in Eq. (4). By this method, we find  $\text{BR}(\Lambda_c^+ \rightarrow pK^+\pi^-)/\text{BR}(\Lambda_c^+ \rightarrow pK^-\pi^+) = (0.05 \pm 0.26 \pm 0.02)\%$  and  $\text{BR}(D_s^+ \rightarrow K^+K^+\pi^-)/\text{BR}(D_s^+ \rightarrow K^-K^+\pi^+) = (0.52 \pm 0.17 \pm 0.11)\%$  where the first errors are statistical and the second are systematic. The BR probability distributions and the 90% limits, as described by Eq. (4), for both DCS decays are shown in Fig. 4. Both limits are larger than the expected ( $\lesssim \tan^4 \theta_c$ ) level, but are the first reported limits on these decays. Furthermore, this is the first successful application of the GP technique to an HEP data analysis.

### Acknowledgements

We wish to acknowledge the assistance of the staffs of Fermi National Accelerator Laboratory, the

INFN of Italy, and the Physics Departments of the Collaborating Institutions. This research was supported in part by the US National Science Foundation, the US Department of Energy, the Italian Istituto Nazionale di Fisica Nucleare and Ministero dell'Istruzione dell'Università e della Ricerca, the Brazilian Conselho Nacional de Desenvolvimento Científico e Tecnológico, CONACyT-México, the Korean Ministry of Education, and the Korean Science and Engineering Foundation.

### References

- [1] H.J. Lipkin, Nucl. Phys. B (Proc. Suppl.) 115 (2003) 117.
- [2] J.R. Koza, Genetic Programming: On the Programming of Computers by Means of Natural Selection, MIT Press, Cambridge, MA, 1992.
- [3] J.M. Link, et al., hep-ex/0503007, Nucl. Instrum. Methods, in press.
- [4] P.L. Frabetti, et al., Nucl. Instrum. Methods A 320 (1992) 519.
- [5] P.L. Frabetti, et al., Nucl. Instrum. Methods A 329 (1993) 62.
- [6] J.M. Link, et al., Nucl. Instrum. Methods A 516 (2003) 364.
- [7] J.M. Link, et al., Nucl. Instrum. Methods A 484 (2002) 174.
- [8] J.M. Link, et al., Nucl. Instrum. Methods A 484 (2002) 270.
- [9] S. Eidelman, et al., Phys. Lett. B 592 (2004) 1.
- [10] M.R. Convery, Incorporating multiplicative systematic errors in branching ratio limits, SLAC-TN-03-001.
- [11] S. Malvezzi, AIP Conf. Proc. 549 (2000) 569.